

On the Leading Order Hadronic Contribution to $(g - 2)_\mu$

Kim Maltman

Math & Stats, York Univ., Toronto, CANADA, and CSSM, Univ. of Adelaide, Adelaide, AUSTRALIA

Abstract. Sum rule constraints dominated by the independent high-scale input, $\alpha_s(M_Z)$, are shown to be satisfied by $I = 1$ spectral data from hadronic τ decays, but violated by the pre-2005 electroproduction (EM) cross-section data. Determinations of the Standard Model (SM) hadronic contribution to $(g - 2)_\mu$ incorporating τ decay data are thus favored over those based solely on EM data, implying a SM prediction for $(g - 2)_\mu$ in agreement with current experimental results.

After accounting for the accurately known, purely leptonic contributions, the largest of the SM contributions to $a_\mu \equiv (g - 2)_\mu/2$ is that due to leading (LO) hadronic vacuum polarization (VP), $[a_\mu]_{had}^{LO}$. This contribution can be expressed as a weighted integral, with known kernel, over the electromagnetic (EM) spectral function $\rho_{EM}(s)$. The uncertainty on $[a_\mu]_{had}^{LO}$ which results is at the $\sim 1\%$ level (comparable to the 0.5 ppm experimental uncertainty on a_μ [1]) and dominates the uncertainty in the SM prediction for a_μ [2]. This determination can, in principle, be improved using CVC, together with data on the $I = 1$ spectral function, $\rho_{I=1}(s)$, obtained from hadronic τ decay. (Isospin-breaking (IB) corrections [3, 4] are needed for accuracy at the 1% level.) Unfortunately, the IB-corrected τ data is in significant disagreement with the corresponding $I = 1$ EM data, $\rho_{I=1}^{EM}$ lying uniformly below $\rho_{I=1}^\tau$ in both the 4π region and that part of the 2π region above the ρ but below 1 GeV^2 [5]. The determination of $[a_\mu]_{had}^{LO}$ employing only EM data leads to a SM a_μ prediction $\sim 2.5\sigma$ below experiment, while the alternate determination incorporating τ decay data is compatible with experiment at the $\sim 1\sigma$ level [5]. We show here that sum rule constraints strongly favor the τ -based determination.

For $w(s)$ any function analytic in the region $|s| < M$, with $M > s_0$, the $I = 1$ vector (V) current and EM correlators, $\Pi(s)$, and corresponding spectral functions, $\rho(s)$, satisfy the FESR relations $\int_0^{s_0} w(s) \rho(s) ds = -\frac{1}{2\pi i} \oint_{|s|=s_0} w(s) \Pi(s) ds$. To suppress duality violation and allow the use of the OPE on the RHS, $w(s)$ should satisfy $w(s = s_0) = 0$ [6]. At scales of $\sim$ a few GeV^2 , the OPE representation for V current correlators is essentially entirely dominated by its leading $D = 0$ component, and hence determined by the single input parameter $\alpha_s(M_Z)$, whose value is known from independent high-scale studies [7]. If one works with weights $w(y)$, $y = s/s_0$, which satisfy $w(1) = 0$, and are non-negative and monotonically decreasing for $0 < y < 1$, the fact that the EM version of $\rho_{I=1}$ lies uniformly below the corresponding τ version implies that

- the normalization *and* slope with respect to s_0 of the EM-based spectral integrals, for a given $w(y)$, should be too low relative to OPE predictions if it is the τ -based

TABLE 1. $\alpha_s(M_Z)$ from fits to the $s_0 = m_\tau^2$ experimental EM and τ spectral integrals, with central $D = 2, 4$ OPE input values

Weight	$[\alpha_s(M_Z)]_{EM}$	$[\alpha_s(M_Z)]_\tau$
w_1	$0.1138^{+0.0030}_{-0.0035}$	$0.1212^{+0.0027}_{-0.0032}$
w_6	$0.1150^{+0.0022}_{-0.0026}$	$0.1195^{+0.0020}_{-0.0022}$

data which is correct and

- the normalization *and* slope with respect to s_0 of the τ -based spectral integrals for a given $w(y)$ should be similarly too high if it is the EM-based data which is correct.

Sum rule tests of the two data sets have been performed for a number of different $w(y)$. See Ref. [8] for details on the OPE and spectral integral inputs (including short- and long-distance IB corrections for the τ data and input relevant to the small $D > 0$ OPE contributions). It is found that both the normalization and slope with respect to s_0 of the τ based spectral integrals are in excellent agreement with OPE expectations, while both the normalization and slope of the EM-based spectral integrals are low, particularly if one uses the pre-2005 EM $\pi\pi$ spectral data.

Table 1 quantifies the EM normalization problem, giving the values of $\alpha_s(M_Z)$ needed to bring the OPE and spectral integrals into agreement for the τ and EM cases, at the maximum common available scale, $s_0 = m_\tau^2$. Results are shown for $w(y) = w_1(y) = 1 - y$ and $w_6(y) = 1 - 6y/5 + y^6/5$ (which have zeros of order 1, 2, respectively, at $s = s_0$). The entries are to be compared, e.g., to the high-scale average, $\alpha_s(M_Z) = 0.1195 \pm 0.0016$, obtained by excluding (i) τ decay input and (ii) the erroneous heavy quarkonium input [9] from the PDG04 average. Table 2 similarly quantifies the EM slope problem. In the OPE column, “indep” and “fit” label results obtained using (i) the independent high-scale $\alpha_s(M_Z)$ input and (ii) the fitted $\alpha_s(M_Z)$ values from Table 1, respectively. We see that lowering the input $\alpha_s(M_Z)$ value to accommodate the EM spectral integral normalizations at $s_0 \sim m_\tau^2$ does *not* resolve the slope problem for the EM spectral integrals.


For illustration purposes, the results for the w_6 EM case are also displayed in Fig. 1. The dotted and solid lines give the central OPE and OPE error bounds, respectively, the solid dots (with error bars) the EM spectral integrals, with pre-2005 $\pi\pi$ input. The open circles show the shifted “EM” spectral integrals obtained by replacing the EM 2π and 4π data with the equivalent τ data. We see that both the EM slope and normalization problems are resolved if, where the $I = 1$ V part of the EM and τ data disagree, the τ data are taken to be correct. The recent 2005 SND EM $\pi\pi$ results [10] are compatible with the corresponding τ results, further strengthening the case for the reliability of the τ data, and the agreement between the SM prediction and experimental result for a_μ .

ACKNOWLEDGMENTS

Thanks to the Natural Sciences and Engineering Research Council of Canada and those authors listed in the acknowledgements of Ref. [8] for their input/assistance.

TABLE 2. Slopes wrt s_0 of the EM OPE and spectral integrals

Weight	S_{exp}	$\alpha_s(M_Z)$	S_{OPE}
w_1	$.00872 \pm .00026$	indep fit	$.00943 \pm .00008$ $.00934 \pm .00008$
w_6	$.00762 \pm .00017$	indep fit	$.00811 \pm .00009$ $.00805 \pm .00009$

**FIGURE 1.** OPE vs. spectral integrals for the w_6 FESR. All notation as described in the text.

REFERENCES

1. G.W. Bennett *et al.* (The Muon $g - 2$ Collaboration), Phys. Rev. Lett. **92**, 161802 (2004) 161802
2. M. Passera, J. Phys. **G31**, R75 (2005)
3. V. Cirigliano, G. Ecker, H. Neufeld, Phys. Lett. **B513**, 36 (2001); JHEP **0208**, 002 (2002)
4. K. Maltman and C.E. Wolfe, hep-ph/0509224, Phys. Rev. **D**, in press
5. M. Davier, S. Eidelman, A. Hocker and Z. Zhang, Eur. Phys. J. **C27**, 497 (2003); Eur. Phys. J. **C31**, 503 (2003). See also other references listed in Ref. [8].
6. K. Maltman, Phys. Lett. **B440**, 367 (1998)
7. QCD review section, S. Eidelman *et al.* (The Particle Data Group), Phys. Lett. **B592**, 1 (2004)
8. K. Maltman, hep-ph/0504201
9. See N. Brambilla *et al.* (The Quarkonium Working Group), hep-ph/0412158, pages 399-400
10. M.N. Achasov *et al.* (The SND Collaboration), hep-ex/0506076